

Veterinary Neutralizer

The neutralizer mixture is added to the pre-enrichment medium whenever the presence of disinfectant solution is known or suspected. As a general rule, this mixture should be used at a rate of 10% of the final volume of the pre-enrichment medium.

► **Storage** : 15 - 25 °C in the dark

Formula in g/l			
Disodium phosphate	85.7	L-Histidine HCL	1
Sodium thiosulphate	5	Tween 80	15
Lecithin	3	Distilled water	890.3

Cat. No.	Description	Units/box	€ Excl. VAT/box
693472	ClearLine®veterinary neutralizer Twist-off bottle 900 ml	9	NC -

Nitrate mobility (Agar)

Medium for confirmation of clostridium perfringens.

► **Storage** : 2 - 25 °C

► **Incubation** : 35 ± 2 °C for 24 - 48 h

Cat. No.	Description	Pack.	€ Excl. VAT
777327	Dehydrated medium nitrate mobility medium - Condalab	500 g	NC -

Harmful bacteria for breweries (Broth)

For the detection of harmful contaminants in brewing.

► **Storage** : 2 - 25 °C

► **Incubation** : 30 - 35 °C for 4 days

Cat. No.	Description	Pack.	€ Excl. VAT
777863	Dehydrated medium for Nocive Brewers bacteria Broth Agar base modified - Condalab	500 g	NC -

Harmful bacteria for breweries (Agar)

For the detection of harmful contaminants in brewing.

► **Storage** : 2 - 25 °C

► **Incubation** : 30 - 35 °C for 4 days

Cat. No.	Description	Pack.	€ Excl. VAT
777862	Dehydrated medium for Nocive Brewers bacteria Agar broth base modified - Condalab	500 g	NC -

Nutritive (Broth)

For the cultivation of non-demanding germs

► **Storage** : 2 - 25 °C

► **Incubation** : 35 ± 2 °C for 18 - 48 h

Cat. No.	Description	Pack.	€ Excl. VAT
777332	Dehydrated medium for nutrient broth - Condalab	500 g	NC -

Nutrient agar

Medium for the cultivation of non-demanding germs in water and clinical samples.

► **Storage** : 2 - 25 °C

► **Incubation** : 35 ± 2 °C for 18 - 24 h

Cat. No.	Description	Pack.	€ Excl. VAT
777328	Dehydrated medium for nutrient agar ISO 6579 / ISO 10273 - Condalab	500 g	NC -

Nutrient agar with sodium chloride ISO 21258

Medium recommended by ISO 21258 for the confirmation of Enterobacteriaceae.

► **Storage** : 2 - 25 °C (Dehydrated base and ready-to-use medium)

► **Incubation** : 37 ± 1 °C for 24 ± 2 h

Dehydrated base

Cat. No.	Description	Pack.	€ Excl. VAT
777331	Dehydrated medium for nutrient agar with sodium chloride ISO 21258 - Condalab	500 g	NC -

Ready-to-use media

Cat. No.	Description	Pack.	€ Excl. VAT
693374	100 ml bottle of nutrient agar with sodium chloride - ClearLine®	10	NC -

O.G.A. (Oxytetracycline Glucose Agar) (O.G.Y.E.)

Recommended for the enumeration of yeasts and moulds in milk, dairy products and food.

► **Storage** : 2 - 25 °C (Dehydrated base and ready-to-use medium)

► **Incubation** : 20 - 25 °C for 5 - 7 days

Dehydrated base

Cat. No.	Description	Pack.	€ Excl. VAT
777335	Dehydrated medium for O.G.A. agar (O.G.Y.E.) - Condalab	500 g	NC -

Ready-to-use media

Cat. No.	Description	Pack.	€ Excl. VAT
693373	100 ml bottle O.G.A. agar (O.G.Y.E.) - ClearLine®	10	NC -

Supplement

For 500 ml of medium.

► **Add 50 mg of oxytetracycline to 5 ml of sterile distilled water**

Cat. No.	Description	Units/box	€ Excl. VAT/box
777791	Oxytetracycline - 50 mg bottle - Condalab	10	NC -

Osmophilic (Agar)

For the detection of osmophilic yeasts in foods

► **Storage** : 2 - 25 °C

► **Incubation** : 20 ± 20 °C for 48 - 72 h

Cat. No.	Description	Pack.	€ Excl. VAT
777337	Dehydrated medium for osmophilic agar - Condalab	500 g	NC -

Ox bile, bacteriological

Ingredient used as a selective inhibitor in culture media such as BLBVB

► **Storage** : 2 - 25 °C

Cat. No.	Description	Pack.	€ Excl. VAT
777448	Ox bile, bacteriological - Condalab	500 g	NC -
778405	Ox bile, bacteriological - Condalab	25 kg	NC -