

TAT (Broth)

For the detection of bacterial contamination. Soy lecithin and Tween® 20 neutralise the conservatives.

➤ **Storage :** 2 - 8 °C

➤ **Incubation :** 35 ± 2 °C for 18 - 48 h

Cat. No.	Description	Units/ box	€ Excl. VAT/ box
994459	9 ml bottle of TAT broth + Tween®20 - ClearLine®	100	NC -
693403	90 ml bottle of TAT broth + Tween®20 - ClearLine®	10	NC -
994460	99 ml bottle of TAT broth + Tween®20 - ClearLine®	10	NC -

T.B.S. (Tryptone bile salt agar) ISO 9308-1

Used for the detection and enumeration of *Escherichia coli* and other coliforms in water by the membrane filtration technique.

➤ **Storage :** 2 - 25 °C

➤ **Incubation :** The membrane is first incubated at 36°C for 2 - 4 hours on TSA agar and then transferred to TBS agar for incubation at 44 ± 0.5°C for 19 - 20 h


Cat. No.	Description	Pack.	€ Excl. VAT
777414	Dehydrated medium TBS agar - Condalab	500 g	NC -

T.B.X. (Tryptone Bile x-Glucuronide - Chromogenic agar)

For the detection and enumeration of *Escherichia coli* in food.

➤ **Storage :** 2 - 25 °C (dehydrated base) and 2 - 8 °C in the absence of light (ready-to-use medium)

➤ **Incubation :** 44 ± 1 °C for 21 ± 3 h



Escherichia coli

Dehydrated base

Cat. No.	Description	Pack.	€ Excl. VAT
777398	Dehydrated medium TBX agar - Condalab	500 g	NC -

Ready-to-use media

Cat. No.	Description	Pack.	€ Excl. VAT
693388	100 ml bottle TBX agar - ClearLine®	10	NC -
693452	200 ml bottle TBX agar - ClearLine®	6	NC -

TCBS

Used for the isolation of *Vibrio cholerae* and *parahaemolyticus* and most other enteropathogenic *Vibrio* species, mainly from fish and seafood.

➤ **Storage :** 2 - 25 °C

➤ **Incubation :** 35 ± 2 °C for 18 - 24 h

Cat. No.	Description	Pack.	€ Excl. VAT
777399	Dehydrated medium TCBS agar - Condalab	500 g	NC -

Tergitol 7

See Chapman TTC

Tetrathionate (Broth)

For the selective enrichment of salmonella in food, water.

➤ **Storage :** 2 - 25 °C

➤ **Incubation :** 35 ± 2 °C for 18 - 24 h

Cat. No.	Description	Pack.	€ Excl. VAT
777400	Dehydrated medium tetrathionate broth - Condalab	500 g	NC -

Supplements

➤ **Add 20 ml of an iodine-iodide solution (6 g of iodine and 5 g of potassium iodide in 20 ml of water) to 10 ml of medium**

Cat. No.	Description	Pack.	€ Excl. VAT
348454-CER	Iodine	100 g	NC -
362405-CER	Potassium iodide	250 g	NC -

Thioglycolate with resazurin (Broth)

Used to detect the presence of viable aerobic, microaerobic, and anaerobic microorganisms in normally sterile products.

➤ **Storage :** 2 - 25 °C (dehydrated base and ready-to-use medium)

➤ **Incubation :** 30 - 35 °C for 3 days

Dehydrated base

Cat. No.	Description	Pack.	€ Excl. VAT
777402	Dehydrated medium thioglycolate fluid medium with resazurin - Condalab	500 g	NC -

Ready-to-use media

Cat. No.	Description	Pack.	€ Excl. VAT
693418	10 ml tube of thioglycolate broth with resazurin - ClearLine®	100	NC -
693389	100 ml bottle of thioglycolate broth with resazurin - ClearLine®	10	NC -

Tryptone Soy Agar (Agar)

Used for the detection and enumeration of *Escherichia coli* and other coliforms in water by the membrane filtration technique according to ISO 9308-1.

➤ **Storage :** 2 - 25 °C

➤ **Incubation :** 36 ± 2 °C for 21 ± 3 h

Cat. No.	Description	Pack.	€ Excl. VAT
777415	Trypticasein Soy Agar (TSA) - Condalab	500 g	NC -

T.S.A. (Trypticasein Soy Agar) (Agar)

A general purpose medium for the growth and isolation of a wide variety of micro-organisms. It can be supplemented with 5-7% sans to determine haemolytic reactions.

➤ **Storage :** 2 - 25°C (dehydrated base) - 2 - 8°C (ready-to-use medium)

➤ **Incubation :** 35 ± 2 °C for 18 - 24 h



Dehydrated base

Cat. No.	Description	Pack.	€ Excl. VAT
777410	Trypticasein Soy Agar (TSA) - Condalab	500 g	NC -

Ready-to-use media

Cat. No.	Description	Pack.	€ Excl. VAT
693390	Trypticase soy agar (TSA) 100 ml bottle - ClearLine®	10	NC -
693455	Trypticase soy agar (TSA) 200 ml bottle - ClearLine®	6	NC -