

## Anaerobic agar

For the culture of anaerobic germs, especially Clostridium.

➤ **Storage 8 - 15 °C**

➤ **Incubation at 35 °C ± 2 °C for 18 - 48 h**

Cat.No.	Description	Pack.	€ Excl. VAT
777174	Dehydrated medium anaerobic agar - Condalab	500 g	NC -

## Baird-Parker Agar

See Baird-Parker agar

## Campylobacter agar

See campylobacter

## Cetrimide agar

See ketrimide agar

## Chapman agar

See chapman agar

## Salmonella chromogenic agar

See Salmonella chromogenic agar

## Brain heart agar

See Heart brain agar

## Columbia agar

See columbia agar

## DRBC agar

See DRBC agar

## DG18 agar

For the detection and enumeration of yeasts and moulds in dry products with an aw < 0.95.

➤ **Storage : 2 - 8 °C in the dark**

➤ **Incubation : 25 °C for 5 days in a humid aerobic atmosphere**

Cat. No.	Description	Pack.	€ Excl. VAT
693356	DG18 Agar 100 ml bottle - ClearLine®	10	NC -

## Actinomycete Isolation agar glycerol

For the isolation and cultivation of actinomycetes in soil and water.

➤ **Storage : 2 - 25 °C**

➤ **Incubation : 30°C ± 2°C for 72 h**

Cat. No.	Description	Pack.	€ Excl. VAT
777816	Dehydrated Actinomycete Isolation Agar Medium - Condalab	500 g	NC -

## Acetate differential agar

To differentiate Shigella from Escherichia coli.

➤ **Storage 8 - 15 °C**

➤ **Incubation at 35 °C ± 2 °C for 7 days**

Cat.No.	Description	Pack.	€ Excl. VAT
777169	Dehydrated medium acetate differential agar Condalab	500 g	NC -

## Middlebrook Agar

See middlebrook agar

## PCA agar

See PCA

## Blood agar

For the determination of haemolytic germs and confirmation of Bacillus cereus and Listeria monocytogenes.

➤ **Storage 8 - 15 °C**

➤ **Incubation at 35 °C ± 2 °C for 18 - 48 h for Listeria monocytogenes**

➤ **Incubation at 30°C for 24 h for Bacillus cereus**

Cat.No.	Description	Pack.	€ Excl. VAT
777823	Blood agar dehydrated medium (ISO7932) - Condalab	500 g	NC -

## Wilson Blair Agar

See Wilson Blair agar

## Giolotti-Cantoni (broth)

Enrichment medium used for the detection of Staphylococcus aureus in foodstuffs.

➤ **Storage : 2 - 25 °C**

➤ **Incubation : 35 ± 2 °C for 40 - 48 h**

➤ **According to ISO 11133 : 37 ± 1 °C for 24 ± 2 - 48 ± 2 h**


Cat. No.	Description	Pack.	€ Excl. VAT
777257	Dehydrated medium Giolotti broth - Cantoni - Condalab	500 g	NC -
777258	Dehydrated medium Giolotti broth - Cantoni - ISO 11133 - Condalab	500 g	NC -

## Hektoen (agar)

Differential selective medium for enteropathogenic bacteria, especially salmonella and shigella. The composition of the medium allows the differentiation of colonies rapidly fermenting one of the 3 sugars (turning from blue to salmon-red) and/or producing H<sub>2</sub>S (black centre).

➤ **Storage : 2 - 25 °C**

➤ **Incubation : 37 ± 1 °C for 20 - 24 h**



Cat. No.	Description	Pack.	€ Excl. VAT
777263	Dehydrated medium Hektoen Agar - Condalab	500 g	NC -

## I.T.C. - Irganon Ticarcillin and potassium Chlorate broth

Broth for selection and enrichment of yersinia enterocolitica.

➤ **Storage : 2 - 25 °C**

➤ **Incubation : 25 ± 1 °C for 44 ± 4 h**

Cat. No.	Description	Pack.	€ Excl. VAT
777265	Dehydrated medium I.T.C. broth - Condalab	500 g	NC -

## Supplement

➤ **To be reconstituted in 8 ml sterile distilled water**

➤ **Storage : 2 - 8 °C**

Cat. No.	Description	Pack.	€ Excl. VAT
777785	Supplement for I.T.C. broth - Condalab	10 bottles for 500 ml	NC -