

## Mac Conkey (broth) - according to pharmacopoeia

For the detection of coliforms in water, milk and pharmaceutical preparations.

➤ **Storage :** 2 - 25 °C (dehydrated base and ready-to-use medium)

➤ **Incubation :** 42 - 44 °C for 24 - 48 h

### Dehydrated base

Cat. No.	Description	Pack.	€ Excl. VAT
777303	Dehydrated medium Mac Conkey Broth - Condalab	500 g	NC -

### Ready-to-use media

Cat. No.	Description	Pack.	€ Excl. VAT
693302	100 ml bottle of Mac Conkey Broth - ClearLine®	10	NC -

## Mac Conkey (agar) - according to pharmacopoeia

Differential medium to guide the identification of lactose-fermenting and non-lactose-fermenting Enterobacteriaceae. It is recommended for the detection of escherichia coli in water, food, dairy products and pharmaceutical preparations.

➤ **Storage :** 2 - 25 °C (dehydrated base and ready-to-use medium)

➤ **Incubation**

➤ **According to ISO 21567 :** 37 ± 1 °C for 20 - 24 h

➤ **According to ISO 21150 :** 30 - 35 °C for 24 h



Enterobacter aerogenes      Escherichia coli

### Dehydrated base

Cat. No.	Description	Pack.	€ Excl. VAT
777298	Dehydrated medium Mac agar Conkey - Condalab	500 g	NC -

### Ready-to-use media

Cat. No.	Description	Pack.	€ Excl. VAT
693369	100 ml bottle of Mac ConkeyBroth* - ClearLine®	10	NC -
693448	100 ml bottle of Mac ConkeyBroth* - ClearLine®	6	NC -

\* Not in accordance with the pharmacopoeia

## Malt (Broth)

Broth for the isolation and culture of yeasts and moulds.

➤ **Storage** 2 - 25 °C

➤ **Incubation :** 30 ± 2 °C for 18 - 72 h

Cat.No.	Description	Pack.	€ Excl. VAT
777305	Dehydrated medium malt extract broth - Condalab	500 g	NC -

## Malt (Agar)

Recommended for the detection, isolation and enumeration of yeasts and moulds.

➤ **Storage :** 2 - 25°C (dehydrated base) - 2 - 8°C (ready-to-use medium)

➤ **Incubation :** 25 - 30 °C for 18 - 72 h

### Dehydrated base

Cat. No.	Description	Pack.	€ Excl. VAT
777304	Dehydrated medium malt extract agar - Condalab	500 g	NC -

### Ready-to-use media

Cat. No.	Description	Pack.	€ Excl. VAT
693437	Slant tube malt extract agar - ClearLine®	100	NC -
693370	100 ml bottle malt extract agar - ClearLine®	10	NC -

## Mannitol mobility nitrate

For the presumptive identification of Enterobacteriaceae based on mannitol fermentation, germ mobility and nitrate to nitrite reduction.

➤ **Storage :** 2 - 25°C (dehydrated base) - 2 - 8°C (ready-to-use medium)

➤ **Incubation :** 35 ± 2 °C for 18 - 24 h

### Dehydrated base

Cat. No.	Description	Pack.	€ Excl. VAT
777306	Dehydrated medium mannitol mobility nitrate - Condalab	500 g	NC -

### Ready-to-use media

Cat. No.	Description	Pack.	€ Excl. VAT
693461	10 ml tube mannitol mobility nitrate - ClearLine®	6	NC -
693412	10 ml tube mannitol mobility nitrate - ClearLine®	100	NC -

## Mannitol salt Agar (MSA)

See Chapman agar

## Marine (Agar)

For the isolation and enumeration of heterotropic marine bacteria.

➤ **Storage :** 2 - 8 °C

➤ **Incubation :** 20 - 25 °C for 24 - 72 h

Cat. No.	Description	Pack.	€ Excl. VAT
777309	Dehydrated medium marine agar medium - Condalab	500 g	NC -

## M-EI (chromogenic agar)

For the detection and enumeration of Enterococcus in water by the membrane filtration method.

➤ **Storage :** 2 - 25 °C

➤ **Incubation :** 41 ± 0.5 °C for 18 - 24 h




Cat. No.	Description	Pack.	€ Excl. VAT
777310	Dehydrated medium for chromogenic agar M-EI - Condalab	500 g	NC -

## Modified M-EI (chromogenic agar)

For the isolation and differentiation of Enterococcus faecalis and Enterococcus faecium

➤ **Storage :** 2 - 8 °C

➤ **Incubation :** 41 ± 0.5 °C for 18 - 24 h



Cat. No.	Description	Pack.	€ Excl. VAT
778436	Dehydrated medium for modified M-EI chromogenic agar - Condalab	500 g	NC -